

Salford PMS Contract Monitoring Audit

INPS



Table of Editions and Contents

| Date | Version | Contents | Output |
|-------------|----------------|-----------------|---------------|
| 12/04/2010 | v1, 29/03/10 | | pdf |
| 21/04/2010 | | | |
| | | | |

Copyright © INPS Ltd 2010

Contents

| | |
|--|----------|
| SALFORD PMS CONTRACT MONITORING AUDIT | 1 |
| Audit criteria | 1 |
| Smoking status | 1 |
| Smoker codes | 1 |
| BMI | 3 |
| Waist circumference | 3 |
| Physical activity status | 3 |
| Carer status | 4 |
| Carers | 4 |
| Blood pressure | 5 |
| Blood pressure and CVD risk | 5 |
| Hypertension and blood pressure | 5 |
| CHD and blood pressure | 6 |
| CHD and cholesterol | 7 |
| Diabetes and HbA1C | 8 |
| Diabetes and blood pressure | 9 |
| Diabetes and cholesterol | 10 |
| COPD | 11 |
| Cervical screening | 12 |

Salford PMS Contract Monitoring Audit

Audit criteria

Smoking status

PMS 1 - Percentage of patients aged 15 and over who have smoking status recorded in the last 27 months (or recorded as never smoked and aged 25 or over)

Practice Population - All active, registered patients

All patients aged 15 and over

Patients aged 15 and over with a latest smoking status recorded except never smoked (see codes below) in the last 27 months

Patients aged 15 and over with NO smoking status (137..) recorded in last 27 months

All patients between 15 and 24 years

Patients between 15 and 24 years with a latest smoking status of never smoked in the last 27 months

Patients between 15 and 24 years with NO smoking status (137..) recorded in last 27 months

All patients over 25 years

All patients over 25 years with a latest smoking status of never smoked in the last 5 years:

1371 Never smoked tobacco

All patients over 25 years with NO smoking status (137..) in last 5 years

Smoker codes

137.. Tobacco consumption

1372. Trivial smoker - <1 cig/day

- 1373. Light smoker - 1-9 cigs/day
- 1374. Moderate smoker - 10-19 cigs/day
- 1375. Heavy smoker - 20-39 cigs/day
- 1376. Very heavy smoker - 40+ cigs/day
- 1377. Ex-trivial smoker (<1/day)
- 1378. Ex-light smoker (1-9/day)
- 1379. Ex-moderate smoker (10-19/day)
- 137A. Ex-heavy smoker (20-39/day)
- 137B. Ex-very heavy smoker (40+/day)
- 137C. Keeps trying to stop smoking
- 137D. Admitted tobacco cons untrue?
- 137F. Ex-smoker - amount unknown
- 137G. Trying to give up smoking
- 137H. Pipe smoker
- 137J Cigar smoker
- 137K. Stopped smoking
- 137L. Current non-smoker
- 137M Rolls own cigarettes
- 137N. Ex-pipe smoker
- 137O. Ex-cigar smoker
- 137P. Cigarette smoker
- 137Q. Smoking started
- 137R. Current smoker
- 137S. Ex smoker
- 137T. Date ceased smoking
- 137V Smoking reduced
- 137X. Cigarette consumption
- 137a. Pipe tobacco consumption
- 137b. Ready to stop smoking
- 137c. Thinking about stopping smoking
- 137d. Not interested in stopping smoking
- 137e. Smoking restarted
- 137f. Reason for restarting smoking
- 137g. Cigarette pack-years
- 137h Minutes from waking to first tobacco consumption

BMI

PMS 2 - Percentage of patients aged 16 and over with height, weight and BMI recorded in the past 3 years

Practice Population - All active, registered patients

All patients aged 16 and over

Patients aged 16 and over with BMI recorded in the last 3 years:

229.. O/E - height

22A.. O/E - weight

22K% Body Mass Index

22K1. Body Mass Index normal K/M2

22K2. Body Mass Index high K/M2

22K3. Body Mass Index low K/M2

22K4. Body mass index 25-29 - overweight

22K5. Body mass index 30+ - obesity

22K6. Body mass index less than 20

22K8 Body mass index 20-24 Normal

Waist circumference

PMS 3 - Waist circumference for those with BMI 25 -29 recorded in the last 3 years

Practice Population - All active, registered patients

All patients with a BMI (Read code 22K) recorded in the last 3 years

All patients with BMI between 25-29 and waist circumference recorded in the last 3 years

22K.. Body Mass Index, and value 25-29

22N0. Waist circumference

Physical activity status

PMS 5 - Percentage of patients aged 16 and over with physical activity status recorded in the last 5 years

Practice Population - All active, registered patients

All patients aged 16 and over

All patients aged 16 and over with a record of physical activity in the last 5 years

13CH. Physical activity

138.. Exercise grading (hierarchy)

Carer status

PMS 7 - Percentage of patients aged 16 and over with carer status recorded. For patients aged up to 50 this need only be recorded once; for patients aged 50 and over this should be recorded in the last 5 years

Practice Population - All active, registered patients

All patients aged between 16 to 49 (inclusive)

All patients aged between 16 to 49 with a carer status recorded:

918A. Carer

918G. Is a carer

9d46. Carer

918r. Not a carer

All patients aged 50 and over

All patients aged 50 and over with a carer status recorded in the last 5 year

Carers

PMS 8 - Percentage of carers identified that have been offered a referral or other support, including provision of information

Practice Population - All active, registered patients

All patients identified as a carer:

918A. Carer

918G. Is a carer

9d46. Carer

All patients identified as a carer who have been offered referral for further support

8HkB. Referral for social services carer's assessment

Blood pressure

PMS 10 - Percentage of patients aged 25 and over with blood pressure recorded in the last 5 years

Practice Population - All active, registered patients

All patients aged 25 and over

All patients aged 25 and over with a blood pressure record in the last 5 years

246.. O/E - blood pressure reading

Blood pressure and CVD risk

PMS 11 - Percentage of patients aged 45 and over with a BP recorded in the last 5 years of 140/90 to 159/99 who have been CVD risk scored

Rationale: all patients with Grade 1 hypertension with risk score >20 should be coded as hypertension and given treatment as per British hypertension society guidelines IV

Practice Population - All active, registered patients

All patients aged 45 and over

All patients aged 45 and over with a BP recorded with BP range 140/90 to 159/99 in the last 5 years

246.. O/E - blood pressure reading

All patients aged 45 and over with a BP recorded with BP range 140/90 to 159/99 in the last 5 years AND also with a CVD risk score:

662k. JBS cardiovascular disease risk <10% over next 10 years

662l. JBS cardiovascular disease risk 10-20% over next 10 years

662m. JBS cardiovascular disease risk >20% over next 10 years

662n. JBS cardiovascular disease risk >20% up to 30% over next 10 years

Hypertension and blood pressure

PMS 12 - Percentage of patients with hypertension whom the last blood pressure is 150/90 or less (measured within the previous 9 months).

For the purposes of this indicator, any patients who have been exception reported, other than recent diagnosis or registration, will be counted as failed to achieve.

Practice Population - All active, registered patients

Same criteria as QOF BP5 but only exceptions allowed are recent diagnosis or registration

Numerator group

Patients with Hypertension from the denominator group below, who have a latest Blood pressure, measured in the last 9 months before 1st April next year, of less than or equal to 150/90

Denominator group

Patients with hypertension:

G2... Hypertensive disease

G20% Essential hypertension, etc

G24..-G2z Secondary hypertension, etc (excluding G24z1)

Who have a latest blood pressure record equal to or less than 150/90 in the 9 months before next 1st April:

246% O/E - blood pressure reading = < 150/90, excluding:

246O. O/E - BP unrecordable

246H. O/E - Arterial pressure index normal

246I. O/E - Arterial pressure index abnormal

246K. Target systolic blood pressure

246L. Target diastolic blood pressure

246M. White coat hypertension

Or patients with hypertension, excluding those with exceptions:

Patients registered in the 9 months before 1st April next year

Or patients with hypertension first diagnosed in the 9 months before 1st April next year

CHD and blood pressure

PMS 13 - Percentage of patients with coronary heart disease in whom the last blood pressure reading is 150/90 or less (measured within the previous 15 months)

For the purposes of this indicator any patients who have been exception reported, other than recent diagnosis or registration, will be counted as failed to achieve.

Practice Population - All active, registered patients

Same criteria as QOF CHD6 but only exceptions allowed are recent diagnosis or registration

Patients with CHD resulting from the denominator group below, who have a latest Blood Pressure entry equal to or less than 150 / 90 recorded in the 15 months before 1st April next year

Denominator group

CHD patients:

G3-G330z (Removed: G331. Prinzmetal's angina, G331.11 Variant angina pectoris, G332 Coronary artery spasm)

G33z.-G3401

G342.-G366.

G38..-G3z..

Gyu3.% [X] Ischaemic heart diseases

who have Numerator data present - a latest blood pressure record equal to or less than 150/90 in the 15 months before 1st April next year;

Or patients with CHD excluding those with exceptions:

Patients registered in the 9 months before 1st April next year

Or patients first diagnosed with CHD in the 9 months before 1st April next year

CHD and cholesterol

PMS 14 - Percentage of patients with coronary heart disease whose last measured cholesterol is 5mmol/l or less (measured in the previous 15 months).

For the purposes of this indicator any patients who have been exception reported, other than recent diagnosis or registration, will be counted as failed to achieve.

Practice Population - All active, registered patients

Same criteria as QOF CHD8 but only exceptions allowed are recent diagnosis or registration

Patients with CHD from the denominator group below, whose latest Cholesterol recorded in any Test SDA entry is equal to or less than 5 mmol/l or less (latest in the 15 months before 1st April next year)

Denominator group

CHD patients (see above)

who have Numerator data present - whose latest Serum cholesterol recorded in any Test SDA entry is equal to or less than 5 mmol/l or less [latest in the 15 months before 1st April next year]:

44OE. Plasma total cholesterol level

44P.. Serum Cholesterol

44P1. Serum Cholesterol normal

44P2. Serum Cholesterol borderline

44P3. Serum Cholesterol raised

44P4. Serum Cholesterol very high

44PH. Total cholesterol measurement

44PJ Serum total cholesterol level

Or patients with CHD, excluding those with exceptions:

Patients registered in the 9 months before 1st April next year

Or patients first diagnosed with CHD in the 9 months before 1st April next year

Diabetes and HbA1C

PMS 15 - Percentage of patients with diabetes in whom the last HbA1c is 8 or less (measured in the previous 15 months).

For the purposes of this indicator any patients who have been exception reported, other than recent diagnosis or registration, will be counted as failed to achieve.

Practice Population - All active, registered patients

All patients aged 17 and over

Same criteria as QOF DM24 but only exceptions allowed are recent diagnosis or registration

Patients aged 17 and over with diabetes resulting from the denominator group below, with a latest Test entry of HbA1c in the 15 months before 1st April next year with a value equal to or less than 8% of HbA1c, HbA1c IFCC standard >64 mmol/mol in last 15 months.

Denominator group

Diabetes patients aged 17 and over:

C10E.% Type 1 diabetes mellitus

C10F.% Type 2 diabetes mellitus (excluding C10F8 Reaven's syndrome)

who have Numerator data present - with a latest Test SDA entry of HbA1C:

42W% Hb. A1C - diabetic control, etc (excluding 42W5. Haemoglobin A1c level - International Federation of Clinical Chemistry and Laboratory Medicine standardised) - see HbA1C IFCC

42c% HbA1 - diabetic control, etc

44TB. Haemoglobin A1c level

44TC. Haemoglobin A1 level

44TL. Total glycosylated Hb level

or HbA1c IFCC:

42W5. Haemoglobin A1c level - International Federation of Clinical Chemistry and Laboratory Medicine standardised

in the 15 months before 1st April next year with a value equal to or less than 8% of HbA1c, or HbA1c IFCC standard ≤ 64 mmol/mol in last 15 months:

Or patients with diabetes, excluding those with exceptions:

Patients who registered in the 9 months before 1st April next year

Patients first diagnosed with diabetes in the 9 months before 1st April next year

Diabetes and blood pressure

PMS 16 - Percentage of patients with diabetes in whom the last BP is 145/85 or less (measured in the previous 15 months).

For the purposes of this indicator any patients who have been exception reported, other than recent diagnosis or registration, will be counted as failed to achieve.

Practice Population - All active, registered patients

All patients aged 17 and over

Same criteria as QOF DM12 but no exceptions (only exceptions allowed are recent diagnosis or registration) - Also date of BP needs to have been measured in the previous 15 months.

Patients aged 17 and over with diabetes, resulting from the denominator group below, who have a latest Blood Pressure record in the 15 months before 1st April next year of less than or equal to 145 / 85

Denominator group

Diabetes patients aged 17 and over:

C10E.% Type 1 diabetes mellitus

C10F.% Type 2 diabetes mellitus (excluding C10F8 Reaven's syndrome)

who have Numerator data present - a latest blood pressure SDA entry in the 15 months before 1st April next year of less than or equal to 145 / 85:

246% O/E - blood pressure reading, excluding:

246O. O/E - BP unrecordable

246H. O/E - Arterial pressure index normal

246I. O/E - Arterial pressure index abnormal

246K. Target systolic blood pressure

246L. Target diastolic blood pressure

246M. White coat hypertension

Or patients with diabetes, excluding those with exceptions:

Patients who registered in the 9 months before 1st April next year

Patients first diagnosed with diabetes in the 9 months before 1st April next year

Diabetes and cholesterol

PMS 17 – Percentage of patients with diabetes whose last measured total cholesterol is 5mmol/l or less (measured in the previous 15 months).

For the purposes of this indicator any patients who have been exception reported, other than recent diagnosis or registration, will be counted as failed to achieve.

Practice Population - All active, registered patients

All patients aged 17 and over

Same criteria as QOF DM17 but only exceptions allowed are recent diagnosis or registration.

Patients aged 17 or over with diabetes, resulting from the denominator group below, whose latest Cholesterol entry recorded in the 15 months before 1st April next year is less than or equal to 5 mmol/l

Denominator group

Diabetes patients aged 17 and over:

C10E.% Type 1 diabetes mellitus

C10F.% Type 2 diabetes mellitus (excluding C10F8 Reaven's syndrome)

who have Numerator data present - whose latest Cholesterol entry recorded in the 15 months before 1st April next year is less than or equal to 5 mmol/l in any test SDA:

44OE. Plasma total cholesterol level

44P.. Serum Cholesterol

44P1. Serum Cholesterol normal

44P2. Serum Cholesterol borderline

44P3. Serum Cholesterol raised

44P4. Serum Cholesterol very high

44PH. Total cholesterol measurement

44PJ Serum total cholesterol level

Or patients with diabetes, excluding those with exceptions:

Patients who registered in the 9 months before 1st April next year

Patients first diagnosed with diabetes in the 9 months before 1st April next year

COPD

PMS 18 – Percentage of patients with COPD who have been categorised as having mild, moderate or severe disease in the last 15 months as defined by FEV1% predicted

Practice Population - All active, registered patients

All patients with COPD:

H3... Chronic obstructive pulmonary disease

All patients with COPD categorised as having mild, moderate, severe disease in the last 15 months:

H36. Mild chronic obstructive pulmonary disease

H37. Moderate chronic obstructive pulmonary disease

H38. Severe chronic obstructive pulmonary disease

with a record of FeV1 in the last 2 years:

3398. FEV1/FVC ratio normal

3399. FEV1/FVC ratio abnormal

339M. FEV1/FVC ratio

339O. Forced expired volume in 1 second

- 339R. FEV1/FVC percent
- 339S. Percent predicted FEV1
- 339T. FEV1/FVC > 70% of predicted
- 339U. FEV1/FVC < 70% of predicted
- 339a. FEV1 before bronchodilation
- 339b. FEV1 after bronchodilation
- 339e. FEV1 pre steroids
- 339f. FEV1 post steroids
- 339j. FEV1/FVC ratio pre steroids
- 339k. FEV1/FVC ratio post steroids
- 339l. FEV1/FVC ratio before bronchodilator
- 339m. FEV1/FVC ratio after bronchodilator

Cervical screening

PMS 19 – Percentage of women aged 25 to 64 that have had cervical screening in the previous 3/5 years as per the KC53 return

Practice Population - All active, registered patients

Percentage of women aged 25 – 49 excluding hysterectomy (EVER) and removed from national programme (last 3 years):

- 685H. No smear - benign hysterectomy
- 685I. No smear - amputation of cervix
- 685K. No smear - no cervix
- 908Y. No cervical smear required - no uterus
- 7E05. Vaginal excision of uterus
- 7E05z Vaginal excision of uterus NOS
- 7E040 Abdominal hysterocolpectomy and excision periuterine tissue
- 7E042 Abdominal hysterocolpectomy NEC
- 7E043 Total abdominal hysterectomy NEC
- 7E046 Radical hysterectomy
- 7E049 TAH – Tot abdom hysterectomy and BSO – bilat salpingophorect
- 7E04B Laparoscopic total abdominal hysterectomy bilat salpingo - oophorectomy
- 7L0A. Clearance of pelvis
- 7L0Az Clearance of pelvis NOS
- 908Q. Cervical Smear disclaimer received – LAST 3 YEARS

Percentage of women aged 25-49 excluding hysterectomy (EVER) and removed from national programme (last 3 years) that have had cervical screening in the previous 3 years

4K22. Cervical Smear negative
4K23. Cervical Smear mild dyskariosis
4K24. Cervical Smear severe dyskariosis
4K25. Cervical Smear severe dysk.? inv. ca
4K26. Cervical Smear: ? gland neopl
4K27. Cervical Smear: atrophic change
4K28. Cervical Smear: mod dyskariosis
4K29. Cervical Smear borderline changes
4K2C. Smear NAD - no endocervical cells
4KA1. Vaginal Vault smear negative
4KA3. Vaginal Vault smear-atrophic
4KA4. Vaginal Vault smear abnormal
6856. Ca Cervix screen - up to date
6859. Ca cervix screen - done
685B. Ca cervix screen - normal
685C. Ca cervix screen - abnormal
685D. Ca cervix screen + fee claim
685R. Liquid based cervical cytology screening
7E2A0 Exam female genital tract under anesth & Papanicolau smear
7E2A2 Papanicolau smear NEC
7E2A3 Vaginal Vault Smear
ZV762 [V] Cervix malignancy screening

Percentage of women aged 50 – 64 excluding hysterectomy (EVER) and removed from national programme (last 5 years)

Percentage of women aged 50-64 excluding hysterectomy (EVER) and removed from national programme (last 5 years) that have had cervical screening in the previous 5 years

NO ALLOWANCE FOR EXCEPTIONS EXCEPT TOTAL HYSTERECTOMIES / REMOVED FROM NATIONAL PROGRAMME
